

Semi-local micro-differential theory and computations of moments of semi-algebraic domains

Gabriela Putinar

Address: Department of Mathematics, University of California, Santa Barbara, CA 93106, U.S.A.

e-mail: gputinar@att.net

Mathematics Sciences Classification (2000). 44A60, 65R32, 14P05, 34M15, 34M35.

Keywords: extremal L -problem of moments, Laplace transform, Gauss-Manin connexion, microdifferential system.

Abstract

In the extremal n -variable L -moment problem, the solution (=the characteristic function of $\{p < 0\}$,) p a polynomial, is determined by finitely many moments, in a set A .

In [23], for quadrature domains ($n = 2$), the reduction of all moments to the moments in A is implicit in the reconstruction of p using hyponormal operators.

For $n \geq 2$, assuming that the complexified of p has only isolated critical points for its singularities, we show that the local algorithm [22] which reduces to the moments in a base extends to the global case, with equality modulo analytic functions. The proofs depend on results from complex singularity theory and micro-differential systems ([15], [20], [21]).

0. INTRODUCTION

0.1. The (n -dimensional, $n \geq 1$) moment problem (in functional analysis) asks for a characterization of those numerical sequences $(a_\alpha)_{\alpha \in \mathbf{N}^n}$ that can be realized as the sequence of *moments* of a positive (Borel) measure μ supported on a subset of \mathbf{R}^n , i.e. s.t.

$$a_\alpha = a_\alpha(\mu) := \int x^\alpha d\mu(x), \quad \alpha \in \mathbf{N}^n.$$

An obvious necessary condition is the non-negativity of the Hankel matrix $(a_{\alpha+\beta})_{\alpha, \beta \in \mathbf{N}^n}$. Versions of this condition are also shown to be sufficient under various conditions on the support of μ (cf. [7], [13], [23] and the references there).

A related problem, the extremal L -moment problem on a fixed compact set in \mathbf{R}^n (with L a bound on the norm of the measures sought, and with extremality

in the set of all moments of such measures), is shown to have a (unique) solution (assuming the Hankel matrix ≥ 0 condition) of the form $\mu = \chi_{\{p < 0\}} dx$ where $p \in \mathbf{R}[x_1, \dots, x_n]$ and dx is the volume form $dx := dx_1 \wedge \dots \wedge dx_n$ (χ denotes the characteristic function of a set).

In the "dynamic" setting, let us consider a family of relatively compact domains $(\Omega_t)_t$, $\Omega_t := \{p < t\}$, t a parameter, associated with a polynomial p , together with their moments $a_\alpha(t)$ with respect to the measure $\chi_{\Omega_t} dx$.

For $n \geq 2$, by the co-area theorem, we have, for t a regular value of p ,

$$a'_\alpha(t) = \int_{\partial\Omega_t} x^\alpha dx/dp,$$

where $\omega := x^\alpha dx/dp$ is a C^∞ $(n-1)$ -form s.t. $dp \wedge \omega = dx$, with uniquely defined restriction to $p^{-1}(t)$.

In this paper we obtain information on real integrals of the form

$$I_f(t) := \int_{\{p=t\}} f(x) dx/dp, \quad (1)$$

(and therefore on $a'_\alpha(t) = a'_{x^\alpha}(t)$), by considering instead complex integrals (in a complex variable t , associated with a complex polynomial -replacing p with its complexified $p_{\mathbf{C}}$ if p is real), of the form

$$I_f^\gamma(t) := \int_{\gamma(t)} f(x) dx/dp. \quad (2)$$

Here the variable $x = (x_1, \dots, x_n)$ is extended to complex values; assuming (*1), section 1.2, $[\gamma(t)] \in H_{n-1}^c(p^{-1}(t))$ is parallel transported (see section 1 for details).

In the local case, such integrals are studied extensively in singularity theory (cf. [1], [15],[19] etc.). For $n = 1$, the question of reduction of integrals to a basis is classical (cf. [8], [19]).

0.2. In the case of an isolated critical point for the complexified $p_{\mathbf{C}}$ of p , by adapting an algorithm (cf. [15]) that expresses n -forms in a certain free $\mathbf{C}[[\tau^{-1}]]$ -module (τ a variable) in terms of a basis, the following local result was proved in [22].

Theorem 0.2.1. *Let $p = p_{\mathbf{C}} \in \mathbf{C}[x]$, $x = (x_1, \dots, x_n)$, be s.t. 0 is a critical point for $p_{\mathbf{C}}$ where $p(0) = 0$. Let $(x^\alpha)_{\alpha \in I}$ be a monomial basis for $\mathbf{C}\{x_1, \dots, x_n\}/I_{\nabla p}$, where $I_{\nabla p}$ is the ideal $(\partial_1 p, \dots, \partial_n p)$. Let $f \in \mathbf{C}[x]$. Then*

a) for every $[\gamma(t)] \in H_{n-1}^c(p^{-1}(t))$ (parallel transported in the homological Milnor fibration),

$$I_f^\gamma = \sum_{\alpha \in I, 0 \leq l \leq k} c_\alpha^l \partial_t^{-l} I_{x^\alpha}, \quad (3)$$

on a (punctured) neighborhood of $0 \in \mathbf{C}$. Here $k = k(f)$ is finite and the coefficients c_α^l are computed by the algorithm of Lemma 0.2.2 below.

b) if $p \in \mathbf{R}[x]$ and if $p \geq 0$ or 1 is not an eigenvalue of the monodromy of $p_{\mathbf{C}}$, we have, asymptotically as $t \rightarrow 0$ ($t > 0$, resp. $t < 0$)

$$a'_f(t) \sim \sum_{\alpha \in I, 0 \leq l \leq k} c_\alpha^l (\partial_t)^{-l} a'_{x^\alpha}(t), \quad (4)$$

mod a power series in t .

We shall denote the expression on the RH of (3) by

$$E_f^\gamma := \sum_{\alpha \in I, 0 \leq l \leq k} c_\alpha^l \partial_t^{-l} I_{x^\alpha}.$$

The notation ∂_t^{-1} will be used below mod \mathcal{O} (cf. section 2). In the above Thm., it is an abbreviation for the algorithm proceeding backwards, from k to 0 , with differentiation of the LH term instead of integration of the RH term (this gives a "continuous fraction" in ∂_t^{-1}).

Lemma 0.2.2. *The coefficients in the expression for E_f are computed starting with $f_0 = f$ by polynomial division and differentiation: for $l \geq 0$,*

$$f_l = \sum_{\alpha \in I} c_\alpha^l x^\alpha + \langle g^l, \nabla p \rangle, \quad f_{l+1} := \operatorname{div} g^l, \quad (5)$$

(here g^l is a vectorial polynomial and $\langle \cdot, \cdot \rangle$ denotes scalar product); the algorithm is finite since $\operatorname{multideg}(f_{l+1}) < \operatorname{multideg} f_l$.

We shall denote $k := \min \{ l \mid f_{l+1} = 0 \}$. The coefficients are unique (Lemma 2 of [PuG], argument using p, f real-valued). If $f \in I_{\nabla p}$ and γ is "generic" (i.e. its orbit under the monodromy group \mathbf{C} -generates $H_{n-1}(X_t)$, X_t = the Milnor fiber) then $I_f^\gamma = 0$ if and only if $f(x)dx/dp$ is an exact form. (This characterizes most complex relations among derivatives of moments.) It is also useful to note for what follows that any writing as in Lemma 0.2.2 for f gives an algorithm as in Thm. 0.2.1 (possibly infinite though).

0.3. In the local case (at 0) one proves, more precisely (19); cf section 2 for definitions) that the \mathcal{O}_t -submodule $\mathcal{K}_p^{(0)}$ of the Gauss-Manin micro-differential module \mathcal{K}_p is free of rank μ over the ring $\mathbf{C}\{\{\partial_t^{-1}\}\}$ of micro-differential operators of order 0 with constant coefficients.

This implies that an $(n-1)$ -form $\omega = f(x)dx/dp$ as above has a unique decomposition

$$f dx/dp = \sum_{\alpha \in I} P_\alpha \cdot x^\alpha dx/dp,$$

where $P_\alpha \in \mathbf{C}\{\{\partial_t^{-1}\}\}$ and $|I| = \text{the Milnor number} = \dim \mathbf{C}\{x\}/(\partial_1 p, \dots, \partial_n p)$.

Since $P_\alpha = \sum_{0 \leq l < \infty} c_\alpha^l (\partial_t)^{-l}$, (with a certain convergence condition on the series of coefficients), integrating, this gives the expression for I_f^γ of Theorem 0.2.1 (but infinite, since $\mathbf{C}[x]$ is replaced with the local ring $\mathbf{C}\{x\}$.)

The assertion on the freeness of $\mathcal{K}_p^{(0)}$ includes therefore the fact that the coefficients are unique, and also the fact that the only relations possible are given by exact forms.

In the τ -variable, multiplication by τ^{-1} in $G_1 := \Omega^n/dp \wedge d(\Omega^{n-2})$ corresponds [Ph0] with the ∂_t^{-1} -action of $\mathbf{C}\{\{\partial_t^{-1}\}\}$ on $\mathcal{K}_p^{(0)}$. An integration by parts argument ([M1]) shows that the algorithm of Thm. 0.2.1 is essentially equivalent with writing in terms of a basis in each of these spaces.

0.4. In the global case, if we assume that p satisfies (*1) (section 1), we can localize using results of [20] and prove (in section 3) the following.

Theorem 0.4.1. *Let $p, f \in \mathbf{C}[x]$, $x = (x_1, \dots, x_n)$, where p satisfies (*1), and let $(x^\alpha)_{\alpha \in I}$ be a monomial basis for $\mathbf{C}[x]/I_{\nabla p}$, where $I_{\nabla p} = \text{the ideal } (\partial_1 p, \dots, \partial_n p) \text{ in } \mathbf{C}[x]$.*

Let $[\gamma(t)] \in H_{n-1}^c(p^{-1}(t))$ be obtained by parallel transport in the cohomology bundle over $\mathbf{C}^ := \mathbf{C} \setminus \mathcal{R}_p$, where $\mathcal{R}_p = \{t_j | 1 \leq j \leq s\} = \text{is the set of critical values of } p$.*

Then, with coefficients c_α^l as in Lemma 0.2.2, we have

a) $I_f^\gamma = E_f^\gamma \pmod{\text{an entire function}}$, for $t \in \mathbf{C}' := \mathbf{C} \setminus \cup_{1 \leq j \leq s} C_j$, where $(C_j)_j$ are parallel cuts at $(t_j)_j$ (in a generic direction θ).

b) if the solutions of the differential system \mathcal{K}_p are of moderate growth at ∞ , then the equality at a) becomes an equality mod a polynomial;

c) if p, f are real-valued and $p \geq 0$, then the moment $a_f(t)$ of $\Omega_t := \{p < t\}$ satisfies

$$a'_f(t) = \sum_{\alpha \in I, 0 \leq l \leq k} c_\alpha^l \partial_t^{-l} a'_\alpha(t)$$

mod a real-analytic function on \mathbf{R} , for $t \gg 0$ (resp. $t \ll 0$).

The set I has cardinality $\mu_T := \text{total Milnor number of } p = \dim_{\mathbf{C}} \mathbf{C}[x]/I_{\nabla p}$ (cf. e.g. [10]).

Uniqueness of the coefficients of the algorithm of Theorem 0.4.1 follows since, by Lemma 3.5, I_f^γ decomposes uniquely as a sum of local $I_f^{e_j}$ mod $\mathcal{O}(\mathbf{C})$. For γ generic and $f \in I_{\nabla p}$, the relations among integrals are characterized by: $I_f^\gamma = 0$ iff for $1 \leq j \leq s$, $f dx/dp$ is exact on a neighborhood of the critical point x_j .

0.5. As to proofs, first, from the local algorithm of Theorem 0.2.1 and a decomposition theorem for n -chains ([20]), the global algorithm via the (Borel)-Laplace transform (in τ) follows immediately (Prop. 3.1).

In order to invert this (Prop. 3.8) to an algorithm in t , injectivity of the semi-local Laplace transform ([21]) under (various) growth conditions ([17]) is needed; with moderate growth conditions, pnt. b) of thm. 0.4.1 follows.

A direct proof (Prop. 3.4) of pnt. a) follows from the local algorithm in t by the variation map ([20]) and a standard local cohomology argument (Lemmas 3.5, 3.6).

In Prop. 3.2 (using [20] -the variation map and (Borel)-Laplace transforms), we prove a version of the local algorithm, for co-vanishing instead of vanishing cycles.

Finally, we apply the above to the real case, to obtain an asymptotics algorithm for oscillatory integrals with phase p (Proposition 3.10); if $p \geq 0$, we obtain as a consequence (Proposition 3.11) the real semi-local algorithm (by the co-area theorem, this proves pnt. c) of Thm. 0.4.1).

These technical results on the semi-local Gauss-Manin are in the spirit of [21], but are different from them in several ways (cf. Remark 3.7). The author could not find any written account of these nor any up-dates in connection with this particular direction. Quite remotely, for extremal moments, the question of reducing moments to a basis was asked in [23] and solved (implicitly, in the reconstruction algorithm for p) for $n = 2$ (in several cases, including that of quadrature domains), by using hyponormal operators.

0.6. The paper is organized as follows: section 1 consists of definitions and properties of the various integrals associated with the polynomial p . In section 2, the complex integrals are viewed as solutions of the Gauss-Manin (micro)-differential system. (These two sections are preliminary). In section 3 we prove several versions of the algorithm of theorem 0.4.1 in the global case. Section 4 consists of examples corroborating the algorithm with asymptotic computations ([4], [18]) and with computations for planar quadrature domains ([2], [12]).

1. INTEGRALS (RELATIVE A POLYNOMIAL p).

1.1. Moments and the Dirac distribution. If $p \geq -M$ (some constant M) is a polynomial in $\mathbf{R}[x] = \mathbf{R}[x_1, \dots, x_n]$, then the moments $a_f(t)$ of $p < t$ are well-defined (for f polynomial) and $I_f(t) = a'_f(t)$ for t a regular value of p (cf. 0.1). (The form fdx/dp in (1) denotes the restriction to $p^{-1}(t)$ of any $(n-1)$ -form η s.t. $dp \wedge \eta = fdx$ - this is independent of the choice of η .)

Let us now consider the local case of domains in a neighborhood of a critical value (0, say) of p . The integral (1) is well-defined if $f \in C_0^\infty$ for p arbitrary in $\mathbf{R}[x]$.

The map from C_0^∞ to \mathbf{R} defined by $f \rightarrow I_f$ is the "Dirac distribution along p ". It has an asymptotic expansion at (the critical value) 0, with coefficients distributions in f , of the form:

$$I_f(t) \sim \sum_{\alpha_0 > -1, 0 \leq k \leq n-1} c_{\alpha_0, k}(f) \cdot t^{\alpha_0} (\log t)^k,$$

where the index α_0 ranges over a sequence of rational numbers (with a single denominator, depending on p only).

If 0 is an isolated critical value for p , then the asymptotic expansion as $t \rightarrow 0$ of the Dirac distribution applied at f depends only on the Taylor series of f at 0; if moreover 0 is an isolated critical value for $p_{\mathbf{C}}$, then the indices α_0, k depend on the monodromy of $p_{\mathbf{C}}$ (cf. [15]).

1.2. Homology decompositions. Let $p : \mathbf{C}^n \rightarrow \mathbf{C}$ be a polynomial. Then p has finitely many branching points (i.e. critical values and "second type" singularities (cf. [20] for the definition).

We shall assume that the following *condition (*1)* is satisfied:

- i) p has no "second type" singularities ([20]); and*
- ii) for every critical value there is only one corresponding critical point.*

We shall denote the (finite) set of critical points (resp. critical values) of p by

$$\mathcal{C}_p = \{x_j \mid 1 \leq j \leq s\}, \text{ resp. } \mathcal{R}_p = \{t_j \mid 1 \leq j \leq s\}. \quad (6)$$

By (*1), p defines a locally trivial fibration over $\mathbf{C}^* := \mathbf{C} \setminus \mathcal{R}_p$. The parallel transport of homology classes is well-defined in \mathbf{C}^* and by [20] we can also localize homology classes, w.r.t. a fixed generic direction of angle θ , as follows.

With notation $S_c^+ = S_c^+(\theta) = \{t \in \mathbf{C} \mid \operatorname{Re} t e^{-i\theta} \geq c\}$, (resp. S_c^- with \geq changed to \leq), letting $\Phi = \Phi(\theta) := \{A \text{ closed } \subset \mathbf{C}^n \mid A \cap p^{-1}(S_c^-) \text{ compact, for some } c > 0\}$, the cohomology groups $H_n^\Phi(\mathbf{C}^n) := \lim_{\leftarrow} H_n(\mathbf{C}^n, p^{-1}(S_c^+))$ are defined ([20]) using semi-algebraic chains. For $c \gg 0$ (sufficient c s.t. all critical values of p are in S_c^-), and if $t \in S_c^+$, the boundary map $\partial : H_n^\Phi(\mathbf{C}^n) \rightarrow H_{n-1}^c(p^{-1}(t))$ is an isomorphism.

We denote by H^c or H homology with compact supports; the coefficients are taken in \mathbf{C} .

By a deformation retract and excision argument, it follows that $H_n^\Phi(\mathbf{C}^n) \cong \bigoplus_{1 \leq j \leq s} H_n(X_j, X_j^+)$, where X_j is a Milnor ball at the critical point x_j and $X_j^+ := p^{-1}(S_j) \cap X_j$, with notation $S_j^+ = t_j + S_\delta^+$ (for a certain small $\delta > 0$.) To this isomorphism corresponds a decomposition

$$\Gamma = \sum_{1 \leq j \leq s} \Gamma_j, \quad (7)$$

for $\Gamma \in H_n^\Phi(\mathbf{C}^n)$.

Applying the boundary map (isomorphism), a similar decomposition holds for $(n-1)$ -cycles. If $\gamma \in H_{n-1}^c(p^{-1}(t))$, then

$$\gamma = \sum_{1 \leq j \leq s} e_j, \quad (8)$$

where e_j is a cycle vanishing in direction θ , i.e. an element in $H_{n-1}(X_{t_{0j}})$, for t_{0j} arbitrary $\in S_j^+$.

1.3. Complex integrals. Let p, f be complex polynomials (n variables), p satisfying (*1). For $[\gamma(t)] \in H_{n-1}^c(p^{-1}(t))$ parallel transported in the cohomology bundle, the complex integrals (2) are well-defined and holomorphic multi-valued for $t \in \mathbf{C} \setminus \mathcal{R}_p$.

From the definitions with forms, if p is real the (complex) integral $I_f^\gamma(t)$ coincides with the (real) integral $I_f(t)$ for t real. This is so for instance if $p \geq 0$, since in this case $\gamma(t) = p_{\mathbf{C}}^{-1}(t) \cap \mathbf{R}^n$ is compact.

For $\Gamma \in H_n^\Phi(\mathbf{C}^n)$, ($\Phi = \Phi(\theta)$ as in 1.2), the integrals with *phase* p and *amplitude* f (both complex polynomials) are well-defined [20] (i.e. convergent; besides obviously holomorphic in τ) by

$$I_f^\Gamma(\tau) := \int_\Gamma e^{-\tau p(z)} f(z) dz, \quad (9)$$

We consider next localized versions at a critical value (0, say) of the above integrals.

Let $X_t =$ the Milnor fibre $:= B \cap p^{-1}(t)$, t in a small disc D s.t. $0 \in D$. Let $H_{n-1}^F(X_t)$ denote cohomology with closed supports; the elements of $H_{n-1}^c(X_t)$ (resp. $H_{n-1}^F(X_t)$) are called vanishing (resp. co-vanishing) cycles.

By replacing, in the above integrals, $\gamma(t)$ with $e(t) \in H_{n-1}^c(X_t)$ (resp. $\epsilon(t) \in H_{n-1}^F(X_t)$), we obtain $I_f^e(t)$ (resp. $I_f^\epsilon(t)$).

These integrals are shown to be holomorphic multi-valued in $D^* = D \setminus 0$ (resp. holomorphic multi-valued in D^* mod holomorphic in D). We shall use the standard notations $\mathcal{O}(D)$ (resp. $\mathcal{O}(D)/\mathcal{O}(D)$) for the set(s) of such functions.

1.4. (Borel)-Laplace transforms. In the local case ($0 =$ critical point for p , $p(0) = 0$), the complex integrals defined above are related as follows [20].

Let $\epsilon = \epsilon_+ \in H_{n-1}^F(X_{t_+})$, $t_+ > 0$ in neighborhood of 0, be a co-vanishing cycle (corresponding to direction $\theta = 0$). The cycle $e_+ = \text{var}(\epsilon) \in H_{n-1}^c(X_{t_+})$ is defined by $\text{var}(\epsilon) = \epsilon' - \epsilon$, where ϵ' is obtained by parallel transporting ϵ along a circuit (counter-clockwise) around 0. This generates an isotopy of the Milnor ball X (s.t. = identity on ∂X) and with it an n -dimensional chain $\Delta_+ \in H_n(X, X^+)$ s.t. $\partial_+ \Delta_+ = e_+$.

Relative to these paths of integration, it is shown (with definitions below) that

$$\int_{\Delta_+} e^{-\tau p} \omega = \mathcal{BL}(I_f^{\epsilon_+})(\tau) = \mathcal{L}(I_f^{\epsilon_+})(\tau), \quad (10)$$

for every holomorphic n -form $\omega = f(z) dz$, with equality of convergent integrals for τ in a sector in \mathbf{C} .

Here \mathcal{BL} denotes the Borel-Laplace transform, well-defined by

$$\mathcal{BL}(\tilde{h})(\tau) := \int_0^\infty e^{-\tau t} \tilde{h} dt,$$

for \tilde{h} holomorphic multi-valued on a sector with origin at 0.

The Laplace transform \mathcal{L} is defined for \tilde{h} as above by

$$\mathcal{L}(\tilde{h}) := \int_{\delta} e^{-\tau t} \tilde{h} dt + \int_{\sigma} e^{-\tau t} \text{var}(\tilde{h}), \quad (11)$$

where σ (resp. δ) is a small circle, oriented clockwise, with base point t_0 near 0 (resp. a half-line in direction $e^{i\theta}$, with origin t_0), and $\text{var}(\tilde{h})$ is the variation map on functions:

$$\text{var}(\tilde{h})(t) = \tilde{h}(e^{-2\pi i} \cdot t) - \tilde{h}(t). \quad (12)$$

2. (MICRO)-DIFFERENTIAL (SEMI)-LOCAL THEORY.

2.1. Solutions. Let $n \geq 1$, $x = (x_1, \dots, x_n)$ be the variable in \mathbf{C}^n and let $p : \mathbf{C}^n \rightarrow \mathbf{C}$ be a polynomial s.t. (*1). For the local study, assume that 0 is an isolated critical point of p of Milnor number μ and that $p(0) = 0$. Let X_t be the Milnor fibre at 0, with t a regular value (near 0) for p .

In the differential case, for $e(t) \in H_{n-1}(X_t)$ a vanishing cycle, the integral $I_f^e(t)$ (cf. 1.3) is a solution (applied at a differential form $f(x)dx/dp$) of the differential system \mathcal{K}_p defined below (the solutions of which are obtained by integrating over slightly more general forms).

Let $\Omega_{x,t/t}^*$ ($* \in \mathbf{N}$), denote the complex of (germs of) relative differential forms (consisting of forms of the type $\sum_{\alpha} a_{\alpha}(x,t)dx^{\alpha}$ with holomorphic coefficients).

Let

$$\Omega_p^* := \Omega_{x,t/t}^* \left[\frac{1}{t-p} \right] / \Omega_{x,t/t}^*$$

be the complex of meromorphic differential forms with pole along the graph of p .

Let \mathcal{D}_t (resp. $\mathcal{D}_{t,x}$) be the ring of (germs at 0) of differential operators in t (resp. (t, x)) with analytic coefficients.

The *differential Gauss-Manin module* is the \mathcal{D}_t -module defined by

$$\mathcal{K}_p := \Omega_p^n / d\Omega_p^{n-1}. \quad (13)$$

The \mathcal{D}_t -structure of \mathcal{K}_p is induced on forms in Ω_p^n from the derivation ∂_t on integrals over the forms. (By the residue map, it coincides with the Gauss-Manin connexion).

The function

$$t \rightarrow I(\zeta)(t) := \int_{e(t)} \text{Res } \zeta$$

has its values in $\tilde{\mathcal{O}}_{t,0}$ (= the space of germs at 0 of multi-valued analytic functions) and the map $\int_{e(t)} : \zeta \rightarrow I(\zeta)$ is a *solution* of \mathcal{K}_p , i.e. an element of $\text{Hom}_{\mathcal{D}_t}(\mathcal{K}_p, \tilde{\mathcal{O}}_{t,0})$.

The homomorphisms $\int_{e(t)}$ as $e(t)$ varies in $H_{n-1}(X_t)$ are all the solutions of \mathcal{K}_p (and in a one-to-one correspondence with the vanishing cycles $e(t)$).

If ζ has a simple pole, writing $\zeta = \omega/2\pi i(p-t)$, with ω a holomorphic n -form, its residue is $\text{Res } \zeta = \omega/dp|_{p^{-1}(t)}$. Writing $\omega = f dx$, with f a holomorphic function, the integral $I_f^\epsilon(t)$ is therefore a solution (applied at ζ) of \mathcal{K}_p . The set of such integrals forms a *lattice* \mathcal{K}_p^0 in \mathcal{K}_p , i.e. an \mathcal{O}_t -module that generates \mathcal{K}_p (freely, with rank μ) as a \mathcal{D}_t -module.

In the micro-differential case, if $\epsilon(t) \in H_{n-1}^F(X_t)$, the integral $I_f^\epsilon(t)$ is a solution (applied at a differential form) of the micro-differential system \mathcal{K}_p .

This system is an \mathcal{E}_t -module with structure extending its \mathcal{D}_t -module structure, where \mathcal{E}_t - the ring of *micro-differential operators* - is a ring containing \mathcal{D}_t in which ∂_t is invertible ($\mathcal{E}_t =$ the micro-localization of \mathcal{D}_t).

The submodule $\mathcal{K}_p^{(0)}$ is free of rank μ over the ring $\mathbf{C}\{\{\partial_t^{-1}\}\}$ of micro-differential operators of order 0 with constant coefficients.

2.2. Semi-local theory. The study of semi-local solutions is described in [21], in the case of (non-characteristic) systems of the form $\mathcal{E}_{t,x}/\mathcal{I}$, where \mathcal{I} is an ideal (t, x are complex variables, $x = (x_1, \dots, x_n)$).

In the case of a single variable t , since \mathcal{E}_t is a p.i.d., it is possible to reduce to a system defined by a single differential polynomial of the form

$$P(t, \partial_t^{-1}) = t^m + \sum_{0 \leq k \leq \infty} c_k(t) \partial_t^{-k},$$

with a certain condition of simultaneous convergence for $c_k \in \mathcal{O}_{t,0}$. Here $m \in \mathbf{N}$ is the *multiplicity* of P .

The space of solutions relative a fixed disk D in \mathbf{C} is defined (for U simply-connected open in D) by

$$\text{Sol}^D(U) = \{u \in \mathcal{O}(U) \mid P_{t_0}(u) \in \mathcal{O}(\overline{D})\},$$

(where the action of ∂_t^{-1} in P_{t_0} is taken by integrating w.r.t. to t_0 , a base point in D - here the convergence conditions ensure this is possible for some D ; the space Sol is independent of the choice of t_0).

The space of "singularities" of solutions is $\text{sol}^D(U) := \text{Sol}^D(U)/\mathcal{O}(\overline{D})$. Both Sol and sol are regarded as sheaves and germs of solutions can be considered. In a semi-local situation one replaces t^m with a small perturbation of it, a polynomial of degree m having all its roots $(t_j)_{1 \leq j \leq m}$ in D . Fixing a generic direction and (parallel) cuts C_j at t_j in that direction, the solutions are uni-valued on

$$D' = D \setminus \cup_{1 \leq j \leq m} C_j. \tag{14}$$

An argument based on index for operators shows that local solutions at a point extend to D' and that $\dim_{\mathbf{C}} \text{sol}^D(D') = m$.

From this, the following decomposition of the space of semi-local solutions (mod \mathcal{O}) is obtained:

$$\text{sol}^D(D') = \bigoplus_{1 \leq j \leq m} \text{sol}^{D_j}(D'_j), \quad (15)$$

where D_j is a small disk at t_j and $D'_j = D_j \setminus C_j$.

Further, the Laplace transform on the space sol is injective and maps the summands of this decomposition to subspaces of $\mathcal{A}^{\leq r}/\mathcal{A}_{\leq -r}$ (growth $\leq Ce^{r|t|}$ resp. $\leq Ce^{-r'|t|}$, where $r = (1/\sqrt{2}) \cdot$ radius of the disk and $r' < r$ is variable).

The (micro-differential) Gauss-Manin module \mathcal{K}_p (discussed at 2.1) has a presentation as an \mathcal{E}_t -module of the form

$$\mathcal{K}_p \cong \mathcal{B}_p / \sum_{1 \leq i \leq n} \partial_{x_i} \mathcal{B}_p,$$

where

$$\mathcal{B}_p := \mathcal{E}_{t,x} / \mathcal{E}_t \cdot (t - p, \partial_{x_i} - \partial_{x_i} p \cdot \partial_t).$$

The module \mathcal{K}_p is not of the form $\mathcal{E}_{t,x}$ considered above and while in principle this case is sufficient for the study of general holonomic systems ([17]), it is possible to obtain (cf. section 3) a similar semi-local decomposition formula by other, simpler means.

3. ALGORITHM FOR MOMENTS - SEMILOCAL / GLOBAL CASE

Let $p \in \mathbf{C}[x] = \mathbf{C}[x_1, \dots, x_n]$, $n \geq 1$, satisfy (*1), let (cf. 1.2) $\mathcal{C}_p, \mathcal{R}_p$ denote its set of critical points (resp. critical values) and let $\theta \in \mathbf{R}$ be the angle of a fixed generic direction in \mathbf{C} .

Proposition 3.1. *Let $p \in \mathbf{C}[x] = \mathbf{C}[x_1, \dots, x_n]$, $n \geq 1$, satisfy (*1) and let $(x^\alpha)_{\alpha \in I}$ be a monomial basis for $\mathbf{C}[x]/I_{\nabla p}$. Let $\Gamma \in H_n^\Phi$, $\Phi = \Phi(\theta)$ (cf. 1.2).*

Then for every $f \in \mathbf{C}[x]$, there exists a finite k (depending on p and f) such that (cf. 1.3 for notations)

$$I_f^\Gamma(\tau) = \sum_{\alpha \in I, 0 \leq l \leq k} c_\alpha^l \left(\frac{1}{i\tau}\right)^l I_\alpha^\Gamma(\tau) \quad (16)$$

with equality asymptotically in $\tau \rightarrow \infty$ (τ in a sector containing a half-line in direction $e^{i\theta}$). The coefficients $c_\alpha^l \in \mathbf{C}$ are computed by the algorithm of Lemma 0.2.2.

PROOF. By [20] (cf. 1.2), Γ has a decomposition $\Gamma = \sum_{1 \leq j \leq s} \Gamma_j$, where Γ_j is admissible, i.e. in $H_n(X^j, X^{+j})$ (where $+$ is w.r.t direction θ). By the local

algorithm (Thm. 0.2.1) in τ for the integral over Γ_j (relative any writing as in Lemma 0.2.2, not necessarily w.r.t. a local basis), we have

$$I_f^{\Gamma_j}(\tau) = \sum_{\alpha \in I, 0 \leq l \leq k} c_\alpha^l \left(\frac{1}{i\tau}\right)^l I_\alpha^{\Gamma_j}(\tau), \quad (17)$$

asymptotically as $\tau \rightarrow \infty$. Noting that the same form $(fdx - \sum_{\alpha \in I, 0 \leq l \leq k} c_\alpha^l x^\alpha dx)$ is integrated at all the critical points of p , and since $I_f^\Gamma = \sum_{1 \leq j \leq s} I_f^{\Gamma_j}$, summing over j in the above formula gives formula (16) for Γ . \square

We have the following algorithm mod \mathcal{O} in the local case, for integrals over *co-vanishing* cycles.

Proposition 3.2. *Let $p, f \in \mathbf{C}[x]$, and let 0 be a critical point for p . Let $\epsilon(t) \in H_{n-1}^F(p^{-1}(t) \cap X)$, $(X \rightarrow D = \text{Milnor fibration at } 0)$ be defined by parallel transportation. Then*

$$I_f^\epsilon(t) = E_f^\epsilon(t) := \sum_{\alpha \in I, 0 \leq l \leq k} c_\alpha^l (\partial_t)_{t_{0+}}^{-l} I_\alpha^\epsilon(t) \text{ mod } \mathcal{O}(D), \quad (18)$$

(multi-valued) for t in $D^* = D \setminus \{0\}$ (and any choice of the base point t_{0+}); the coefficients $c_\alpha^l \in \mathbf{C}$ are computed by the algorithm of Lemma 0.2.2.

PROOF. By the local algorithm (cf. 0.2.1) for $e := \text{var}(\epsilon)$ (a vanishing cycle), we have $I_f^\epsilon(t) = E_f^\epsilon(t)$ for $t \in D^*$. Therefore also $\mathcal{BL}(I_f^\epsilon)(\tau) = \mathcal{BL}(E_f^\epsilon)(\tau)$ for τ in an infinite sector. By [20] (cf. 1.4), this is equivalent with $\mathcal{L}(I_f^\epsilon)(\tau) = \mathcal{L}(E_f^\epsilon)(\tau)$, from which it follows that $I_f^\epsilon(t) = E_f^\epsilon(t) \text{ mod } \mathcal{O}(D)$, since the local Laplace transform is an injection on $\hat{O}(S)/\hat{O}(D)$ (cf. [17], [21]), where S is a sector in D . \square

Notations 3.3. (In \mathbf{C} , w.r.t. \mathcal{R}_p and a fixed θ .) For $R > 0$, let $S = S_R := \{|t| < R \mid \text{Re}(te^{-i\theta}) < 0\} \cup \{t \mid \text{Re}(te^{-i\theta}) \geq 0 \text{ and } |\text{Im}(te^{-i\theta})| < R\}$ (i.e. a half-disk \cup a half-strip), let $C_j = \{\text{Re}(t - t_j)e^{-i\theta} \geq 0\}$ be parallel cuts at t_j . Let S_j ($1 \leq j \leq s$) be open "thickenings" of C_j containing t_j . Then $C_j \subset S_j \subset S$, for $1 \leq j \leq s$.

Let $S' := S \setminus \cup_{1 \leq j \leq s} C_j$ and $S'_j := S_j \setminus C_j$; $\mathbf{C}' := \mathbf{C} \setminus \cup_{1 \leq j \leq s} C_j$, $\mathbf{C}^* = \mathbf{C} \setminus \mathcal{R}_p$; similarly D', D^* , for D a disk.

Proposition 3.4. *Let $p, f \in \mathbf{C}[x]$, where p satisfies (*1) and let $\gamma = [\gamma(t)] \in H_{n-1}^c(p^{-1}(t))$ be defined by parallel transportation, univalued for $t \in \mathbf{C}'$ (cf. 3.3, w.r.t. a direction θ). Then*

$$I_f^\gamma(t) = E_f^\gamma(t) := \sum_{\alpha \in I, 0 \leq l \leq k} c_\alpha^l (\partial_t)^{-l} I_\alpha^\gamma(t) \text{ mod } \mathcal{O}(\mathbf{C}'), \quad (19)$$

(uni-valued) for t in \mathbf{C}' ; the coefficients $c_\alpha^l \in \mathbf{C}$ are computed by the algorithm of Lemma 0.2.2.

PROOF. a) Since by (*1), p defines a locally trivial fibration over \mathbf{C}' , $[\gamma(t)]$ can be transported by parallelism to $[\gamma(t_0)]$, $t_0 \in S_c^+(\theta)$, $c \gg 0$.

By [PhI], $[\gamma(t_0)] = \sum_{1 \leq j \leq s} [e_j]$, $e_j \in H_{n-1}^c(X_{t_0j})$, vanishing cycles at t_j (t_{0j} in X_j^+ , $+$ being w.r.t. direction θ).

b) By the local algorithm (Thm. 0.2.1) w.r.t. direction θ , $I_f^{e_j}(t) = E_f^{e_j}(t)$, for $1 \leq j \leq s$, t in X_j^+ .

c) For $S = S_R$ (notations as in 3.3) there is a (natural) decomposition (cf. Lemma 3.5 below)

$$\mathcal{O}(S')/\mathcal{O}(S) = \bigoplus_{1 \leq j \leq s} \mathcal{O}(S'_j)/\mathcal{O}(S_j).$$

For $t \in S'$, by a) and Lemma 3.6 below

$$I_f^\gamma(t) = \sum_{1 \leq j \leq s} I_f^{e_j}(t) \bmod \mathcal{O}(S)$$

and similarly

$$E_f^\gamma(t) = \sum_{1 \leq j \leq s} E_f^{e_j}(t) \bmod \mathcal{O}(S).$$

Using b), it follows that $\sum_{1 \leq j \leq s} I_f^{e_j}(t) = \sum_{1 \leq j \leq s} E_f^{e_j}(t) \bmod \mathcal{O}(S)$ for $t \in S'$. Therefore also, $I_f^\gamma(t) = E_f^\gamma(t) \bmod \mathcal{O}(S)$, for $t \in S'$. Letting $R \rightarrow \infty$, since solutions extend to \mathbf{C}^* , we have equality mod $\mathcal{O}(\mathbf{C})$. \square

Using (8), from Lemmas 3.5 and 3.6 below it will follow that the integral $I_{f(z)}^\gamma(t)$ over a global cycle $[\gamma] \in H_{n-1}(p^{-1}(t))$, $t \in S_c^+(\theta)$, $c \gg 0$, localizes, i.e.

$$I_f^\gamma(t) = \sum_{1 \leq j \leq s} I_f^{e_j}(t) \bmod \mathcal{O}(\mathbf{C}).$$

We prove this by using standard arguments from local cohomology.

Lemma 3.5. *With notations as in 3.2, we have*

$$\mathcal{O}(S')/\mathcal{O}(S) \cong \bigoplus_{1 \leq j \leq s} \mathcal{O}(S'_j)/\mathcal{O}(S_j) \tag{20}$$

PROOF. Since $Z := \cup_{1 \leq j \leq s} C_j$ is closed in S , we have the following exact sequence in local cohomology with supports in Z ([9], cf. also [3]).

$$0 \rightarrow H_Z^0(S, \mathcal{O}) \rightarrow H^0(S, \mathcal{O}) \rightarrow H^0(S \setminus Z, \mathcal{O}|_{S \setminus Z}) \rightarrow H_Z^1(S, \mathcal{O}) \rightarrow H^1(S, \mathcal{O}) \rightarrow \dots$$

The leftmost term =0, by the identity principle for (one variable) holomorphic maps; the rightmost term =0 since S (= an open domain in \mathbf{C}) is Stein. Hence the exact sequence

$$0 \rightarrow \mathcal{O}(S) \rightarrow \mathcal{O}(S') \rightarrow H_Z^1(S, \mathcal{O}) \rightarrow 0,$$

or equivalently

$$\mathcal{O}(S')/\mathcal{O}(S) \cong H_Z^1(S, \mathcal{O}).$$

Similarly,

$$\mathcal{O}(S'_j)/\mathcal{O}(S_j) \cong H_{C_j}^1(S_j, \mathcal{O}|_{S_j}).$$

On the other hand, by excision for local cohomology one has

$$H_Z^1(S, \mathcal{O}) \cong \bigoplus_{1 \leq j \leq s} H_{C_j}^1(S, \mathcal{O})$$

and since C_j is closed in S_j and S_j is open in S , also

$$H_{C_j}^1(S, \mathcal{O}) \cong H_{C_j}^1(S_j, \mathcal{O}|_{S_j}).$$

From this the lemma follows immediately. \square

Lemma 3.6. *Let p satisfy (*1) and let S (cf. notations 3.3) contain \mathcal{R}_f . Let $[\gamma] = \sum_{1 \leq j \leq s} [e_j]$ in $H_{n-1}^c(p^{-1}(t_o)) \cong \bigoplus_{1 \leq j \leq s} H_{n-1}^c(X_{t_j}^+)$.*

Then $I_f^\gamma(t) = \sum_{1 \leq j \leq s} I_f^{e_j}(t) \bmod \mathcal{O}(S)$ uni-valued for $t \in S'$, with equality meaning corresponding elements via the isomorphism of Lemma 3.5.

PROOF. Since by (*1) the variation map is well-defined on S , after parallel transporting $[\gamma]$ to S , we have that $\text{var}(\gamma) = \sum_{1 \leq j \leq s} \text{var } e_j$; since the map var on functions commutes with var on cycles [15], [14], it follows that $\text{var } I_f^\gamma = \sum_{1 \leq j \leq s} \text{var } I_f^{e_j}$, as cocycles in $H_Z^1(S, \mathcal{O})$, where $Z := \bigcup_{1 \leq j \leq s} C_j$. Since by excision $H_Z^1(S, \mathcal{O}) \cong \bigoplus_{1 \leq j \leq s} H_{C_j}^1(S, \mathcal{O})$, the same argument as in Lemma 3.5 finishes the proof, since $\text{var} = \delta : H^0(S, \mathcal{O}) \rightarrow H_Z^1(S, \mathcal{O})$ (the Čech coboundary map associates to a section the corresponding cocycle). \square

Remark 3.7. We have essentially avoided using holonomic differential systems, by ad-hoc arguments for the case of the Gauss-Manin differential system; this is a prototype for some of the general theory, cf. [17], [20], [21].

The local Laplace transform (at 0) in direction $a = e^{i\theta}$ is described as follows. Let

$$\mathcal{Z}_a^- := \{Z \mid Z \text{ closed homogeneous } \subset \mathbf{C} \text{ s.t. } \langle z, a \rangle < 0, z \in Z \setminus \{0\}\}.$$

The Laplace transform is the map

$$\mathcal{L}^- = (\mathcal{L}^-)_a : \lim_{r \rightarrow 0} \{Z \in \mathcal{Z}_a^-, r \rightarrow 0\} \mathcal{O}(D_r \setminus Z) / \mathcal{O}(D_r) \longrightarrow (\mathcal{B}^{<1} / \mathcal{B}^{\leq -1})_a$$

(cf. notation below for the target space) defined on $h = h(t)$ s.t. $h = \text{var}(\tilde{h})$ (surjectivity of var , cf. [M0]) by

$$\mathcal{L}^-(h)(\tau) := \int_{\sigma_0} e^{-\tau t} \tilde{h}(t) dt + \int_{\delta} e^{-\tau t} h(t) dt,$$

where $\sigma_0 = \partial D_r$ (based at t_0) and δ is a half-line with origin t_0 in direction a .

By [17], the (local) Laplace transform is an isomorphism between the following spaces ai) and bi), $i = 1, 2, 3$ of holomorphic functions on sectors:

a1) no growth conditions (=: $\text{sp } \mathcal{O}$, specialization of Sato); a2) sub-exponential growth as $t \rightarrow \infty$ (=: $\mathcal{B}^{<1}$); a3) moderate growth as $t \rightarrow \infty$ (=: $\mathcal{B}^{\leq 0}$);

b1) $\mathcal{B}^{<1}/\mathcal{B}^{\leq -1}$; b2) $\mathcal{B}^{<1}$; b3) sub-exponential growth as $\tau \rightarrow \infty$, moderate growth as $\tau \rightarrow 0$ (=: $\mathcal{B}^{<1}$).

In the semi-local case, $\mathcal{L}_{s.l.}^-$ is defined by ([PhII])

$$\mathcal{L}_{s.l.}^-(h)(\tau) := \int_{\sigma} e^{-\tau t} \tilde{h}(t) dt + \sum_{1 \leq j \leq s} \int_{\delta_j} e^{-\tau t} h(t) dt,$$

where $h = \text{var } \tilde{h}$, σ is s.t. $[\sigma] = \sum_{1 \leq j \leq s} [\sigma_j]$ (σ_j and δ_j as in the local case above).

Therefore $\mathcal{L}_{s.l.}^-(h)(\tau) = \sum_{1 \leq j \leq s} \mathcal{L}_{s.l.}^-(h_j)(\tau)$, if $h = \sum_{1 \leq j \leq s} h_j$ (in the sense of the isomorphism of lemma 3.5). Replacing in the spaces bi) above < 1 with ≤ 1 , we obtain spaces bi'), which are codomains for the semi-local Laplace transform.

Proposition 3.8. *Let p, f, γ and a direction $a = e^{i\theta}$ be as in prop. 3.4; let $[\gamma] = \sum_{1 \leq j \leq s} [e_j]$ (as in 1.2).*

If, for $1 \leq j \leq s$, $\mathcal{L}_{s.l.}^-(I_f^{e_j})$ and $(\mathcal{L}_{s.l.}^-(I_{x^\alpha}^{e_j}))_{\alpha \in I}$ are in the space b1) (resp. b2), resp. b3), then $I_f^\gamma = E_f^\gamma$ mod a function which is entire (resp. entire of sub-exponential growth at infinity, resp. a polynomial).

PROOF. 1) In the local case (at t_j), since (by Thm. 0.2.1) $I_f^{e_j} = E_f^{e_j}$ (on D'_j) mod $\mathcal{O}(D_j)$, we have $\mathcal{L}_a^-(I_f^{e_j})(\tau) = \mathcal{L}_a^-(E_f^{e_j})(\tau)$ in $\mathcal{B}^{<1}/\mathcal{B}^{\leq -1}(\Sigma')$, for some (infinite) sector Σ'_j containing direction a . Since \mathcal{L}_a^- is an isomorphism ([17]) between ai) and bi) ($i=1,2,3$), the assertions in the local case follow (with $\mathcal{O}(\mathbb{C})$ replaced with $\mathcal{O}(D_j)$).

2) In the semi-local case ($S = S_R$, R fixed, sufficiently large) since, as in the proof of prop. 3.4, $I_f^\gamma = \sum I_f^{e_j}$ and $E_f^\gamma = \sum E_f^{e_j}$ mod $\mathcal{O}(S)$, with all terms defined on S' (resp. S'_j), taking their semi-local Laplace transforms (using additivity and a commutative square with equalities mod $\mathcal{O}(S)$), we have that $\mathcal{L}_{s.l.}^-(I_f^\gamma)(\tau) = \mathcal{L}_{s.l.}^-(E_f^\gamma)(\tau)$ for $\tau \in \Sigma' \subset \cap \Sigma_j$, for Σ' a sector, with equality in the space b1'.

3) The assertions of the statement of the proposition are now consequences of 2), extendability (by parallel transport of $[\gamma]$) of I_f^γ , E_f^γ , and the fact that

$\mathcal{L}_{s,l}^-$ is injective on ai) $1 \leq i \leq 3$ ([21], the inverse in the semi-local case also given by integration (\mathcal{L}^+ , cf. [17]). \square

Remarks 3.9. 1) In the proof of Proposition 3.4, we can avoid the local cohomology argument of lemmas 3.5 and 3.6 by using proposition 3.1 and inversion of $\mathcal{L}_{s,l}^-$; the solutions I_f^γ and E_f^γ extend analytically over S_R (by parallel transport of $[\gamma]$), therefore $\mathcal{L}_{s,l}^-$ is defined and injective [21] (with target space, see 2.2, $\mathcal{A}^{\leq r}/\mathcal{A}_{\leq -r}$).

2) In the local situation, instead of the spaces in [21], we can use the space \mathcal{A} of functions holomorphic on sectors truncated to a neighborhood of ∞ . Then, since the local Laplace transform is an isomorphism [17] to $\mathcal{B}^{\leq 1}/\mathcal{B}^{< -1}$, we obtain the equivalence between the local algorithms in t and in τ .

Proposition 3.10. *Let $p, f \in \mathbf{R}[x_1, \dots, x_n]$. Then (with coefficients computed as in Lemma 0.2.2),*

$$\int_{\mathbf{R}^n} e^{i\tau p(x)} f(x) dx \sim \sum_{\alpha \in I, 0 \leq k \leq l} c_\alpha^l(i\tau)^{-l} \int_{\mathbf{R}^n} e^{i\tau p(x)} x^\alpha dx,$$

with \sim meaning asymptotically as $\tau \rightarrow \pm\infty$.

PROOF. a) The (phase) integrals are well-defined by truncating f with a C_0^∞ -function which is identically one in an open set containing all critical points of $p_{\mathbf{C}}$, and = 0 outside a neighborhood of its closure, since (e.g. by [15]) such integrals are asymptotically 0 if $\text{supp } f \cap \mathcal{C}_p = \emptyset$.

b) By a partition of unity, it is therefore sufficient to prove the local asymptotic equality:

$$\int_{\mathbf{R}^n} e^{i\tau p(x)} f(x) \phi_j(x) dx \sim \sum_{\alpha \in I, 0 \leq l \leq k} c_\alpha^l(i\tau)^{-l} \int_{\mathbf{R}^n} e^{i\tau p(x)} x^\alpha \phi_j(x) dx,$$

where ϕ_j is a C_0^∞ -function which is identically one in a neighborhood of x_j and = 0 outside a larger neighborhood. We may, by a change of coordinates, assume $x_j = 0$, ($\phi = \phi_j$).

c) At 0, by a theorem of [15], there exists an admissible n -chain $\Gamma = \Gamma^+$ (i.e. in $H_n(X, X^+)$, where $X^+ = \{ \text{Im } p_{\mathbf{C}} > 0 \}$) such that

$$\int_{\mathbf{R}^n} e^{i\tau p(x)} f(x) \phi(x) dx \sim \int_{\Gamma} e^{i\tau p(z)} f(z) dz,$$

asymptotically as $\tau \rightarrow \infty$ (and similarly w.r.t. a Γ^- and $\tau \rightarrow -\infty$.)

In the same way, (for the same n -chains)

$$\sum_{\alpha \in I, 0 \leq k \leq l} c_\alpha^l(i\tau)^{-l} \int_{\mathbf{R}^n} e^{i\tau p(x)} x^\alpha \phi(x) dx \sim \sum_{\alpha \in I, 0 \leq l \leq k} c_\alpha^l(i\tau)^{-l} \int_{\Gamma} e^{i\tau p(z)} z^\alpha dz,$$

asymptotically as $\tau \rightarrow \infty$ (resp. as $\tau \rightarrow -\infty$.)

d) By the above, we have reduced the proof to showing

$$\int_{\Gamma} e^{i\tau p(z)} f(z) dz \sim \sum_{\alpha, l} c_{\alpha}^l (i\tau)^{-l} \int_{\Gamma} e^{i\tau p(z)} z^{\alpha} dz,$$

asymptotically as $\tau \rightarrow \infty$ (resp. as $\tau \rightarrow -\infty$.) which is the local algorithm (thm. 0.2.1), via the Borel-Laplace transform. \square

Proposition 3.11. *Let $p \geq 0$, $p \in \mathbf{R}[x_1, \dots, x_n]$ be s.t. its complexified $p_{\mathbf{C}}$ satisfies (*1), let $\mathcal{R} = \mathcal{R}_p \cap \mathbf{R}$ and let $f \in \mathbf{R}[x_1, \dots, x_n]$.*

Then $\gamma(t) := \{p = t\}$ defines a class in $H_{n-1}^c(p^{-1}(t))$ s.t. the real integrals (cf. 0.1) $I_f(t)$ and $E_f(t)$ (the latter mod \mathcal{O}) are real-analytic on $t > t_+ := \sup \mathcal{R}$ (resp. $t < t_- := \inf \mathcal{R}$).

For $t > t_+$ (resp. $t < t_-$) they satisfy

$$I_f(t) = E_f(t) \tag{21}$$

mod a real-analytic function on \mathbf{R} .

Moreover (writing $[\gamma] = \sum_j [e_j]$ as before), if the complex integrals $I_f^{e_j}$ and $(I_{x_{\alpha}}^{e_j})_{\alpha \in I}$ are of moderate growth at infinity, then the above equality holds mod a (real-valued) polynomial.

The last condition above holds if all the local solutions of the Gauss-Manin system are of moderate growth at infinity.

PROOF. (in the case $t > t_+$, the case $t < t_-$ is similar).

a) By (*1) for p , $[\gamma(t)]$ is well-defined by parallel transportation.

b) By [20], if $c \gg 0$ (sufficient, cf. 1.2, c s.t. all critical values of p are in S_c^-), for $t \in S_c^+$ we have

$$H_{n-1}^c(p^{-1}(t)) \cong H_n(\mathbf{C}^n, p^{-1}(S_c^+)) \cong H_n^{\Phi}(\mathbf{C}^n).$$

Here $S^+ = S^+(\theta)$, $\Phi = \Phi(\theta)$, where θ is a generic direction, arbitrary for now.

c) It is possible to choose θ s.t. the cuts $C_j = C_j(\theta)$ at t_j (abusively, with the same indexing for points in \mathcal{R} as for \mathcal{R}_p) in direction θ satisfy

1) S_c^+ intersects the semi-axis $t > 0$ (e.g. if $\theta \neq -\pi/2$);

2) $C_j \cap \mathbf{R} = \mathcal{R}$ (e.g. if $\theta \neq 0$);

3) θ generic in neighborhood of 0 s.t. $(C_j)_j$ are disjoint.

Then (letting $\theta \rightarrow 0$) a sector Σ_+ with origin at t_+ and axis $\{t > t_+\}$ is in the domain of definition of $I_f^{\gamma}(t)$ and $E_f^{\gamma}(t)$; at the same time, for t in this range, γ admits a decomposition $[\gamma] = \sum [e_j]$ as before (1.2).

d) By Proposition 3.4 for γ , $I_f^{\gamma}(t) = E_f^{\gamma}(t) \bmod \mathcal{O}(\mathbf{C})$, for $t \in \mathbf{C}^*$, equality of univalued holomorphic functions on sectors avoiding $C_j, 1 \leq j \leq s$, in particular on Σ_+ .

For $t \in \Sigma_+$, t real (i.e. $t > t_+$), we have by definition $I_f^\gamma(t) = I_f(t)$ and similarly (since defined mod $\mathcal{O}(\mathbf{C})$, univalued) $E_f^\gamma(t) = E_f(t)$. Therefore for $t \in \Sigma_+$, $I_f(t) - E_f(t) = h(t)$, where $h \in \mathcal{O}(\mathbf{C})$; since it is real-valued, h is real-analytic. This proves (21).

e) The last assertion of the proposition follows from Proposition 3.8 (moderate growth conditions and the Laplace transform). \square

Remark 3.12. For p real ($p \geq 0$) we have used complex integrals over the real cycle $\gamma(t) = \{p = t\}$ to prove an algorithm between real integrals (= derivative of moments associated with p). Another cycle, γ_1 (isotopic with γ) is used for computing local (at 0 = critical point for $p_{\mathbf{C}}$) real asymptotics using complex integrals over γ_1 . The latter satisfies $\gamma_1(t) \subset \{\text{Im } p_{\mathbf{C}} = t\}$, therefore $\gamma_1(t) \cap \mathbf{R}^n = \emptyset$, for any $t > 0$, in particular $\gamma \neq \gamma_1$. Indeed, computations for asymptotics at a critical point by the method of the steepest descent, use directions for which $\text{Im } p_{\mathbf{C}} = \text{const.}$ ($p_{\mathbf{C}}$ = the phase).

In conclusion, we have

PROOF OF THM. 0.4.1. a) follows from prop. 3.4; b) from the fact that I_f^ε and $I_{x^\alpha}^\varepsilon$ are solutions of \mathcal{K}_p (see 2.1) and by prop. 3.8; c) is prop. 3.11.

4. EXAMPLES

We shall abbreviate I_f^γ as I_f and mod $\mathcal{O}(\mathbf{C})$ as mod \mathcal{O} .

4.1. Let $n = 2$ and let

$$p(x, y) = p_\lambda(x, y) = \frac{x^3}{3} - \lambda^2 x + y^2, \quad (22)$$

where $\lambda \in \mathbf{C}$ is a parameter, $x, y \in \mathbf{C}$ are the variables. (This is the mini-versal deformation of the A_2 singularity).

Since $\nabla p = (x^2 - \lambda^2, y)$, a monomial \mathbf{C} -basis for $\mathbf{C}[x, y]/I_{\nabla p}$ is $\{1, x\}$. For $\lambda \neq 0$ there are two non-degenerate critical points $(\pm\lambda, 0)$, which for $\lambda = 0$ degenerate to a single critical point (0, of Milnor number = 2).

In general $I_p = t \cdot I_0$; we have, for this p ,

$$\frac{1}{3}I_{x^3} - \lambda^2 I_x + I_{y^2} = t \cdot I_0. \quad (23)$$

We shall apply the algorithm of section 3 (which shows that I_f can be reduced mod \mathcal{O} to integrals (I_{x^α}) in a basis with indices $\alpha \in I$, $|I| = \mu_T$) to $f = x^3$ and to $f = y^2$.

For $f = x^3$, since $f = f_0 = x(x^2 - \lambda^2) + x\lambda^2$, we have $f_0 = x\lambda^2 + x\partial_x p$, therefore $g^0 = (x, 0)$ and $f_1 = \text{div } g^0 = 1$. Hence

$$I_{x^3} = \lambda^2 I_x + \partial_t^{-1} I_0 \text{ mod } \mathcal{O}. \quad (24)$$

Similarly, from $f_0 := y^2 = (1/2)y\partial_y p$, we have

$$I_{y^2} = \frac{1}{2}\partial_t^{-1}I_0 \bmod \mathcal{O}. \quad (25)$$

Consequences:

a) These latter formulas reduce (23) to a (micro-)differential equation in I_x (if I_0 is known):

$$2\lambda^2 I_x = \left(\frac{5}{2}\partial_t^{-1} - 3t\right)I_0 \bmod \mathcal{O}.$$

For $\lambda = 0$, this equation has solutions $I_0 = c t^{-\frac{1}{6}}$, c a constant; comparing, the Picard-Fuchs equations (at the only critical point 0) give the same solutions for I_0 . Indeed, note that since p_0 is quasi-homogeneous (of quasi-homogeneity degree = 1), the monodromy matrix is $M = \text{diag} (\langle \nu, \alpha + 1 \rangle - 1)_{\alpha \in I}$, where $\nu = (1/3, 1/2)$ is the weight of quasi-homogeneity.

b) We may alternatively reduce (23) to an equation for I_{x^3} in terms of I_0 :

$$-\frac{2}{3}I_{x^3} = \left(t - \frac{3}{2}\partial_t^{-1}\right)I_0.$$

For $\lambda = 0$, this and the equation for I_0 (at *a*) above) imply

$$I_{x^3} = \partial_t^{-1}I_0.$$

If $\gamma(t) = \partial\Omega_t$, (Ω_t relatively compact in \mathbf{R}^n) this follows from

$$I_{x^3} = \int_{\gamma(t)} \frac{x^3 dx dy}{dp} = \int_{\gamma(t)} x dy = \int_{\Omega_t} d(x dy) = \text{vol}(\Omega_t),$$

the last term being $= \partial_t^{-1}I_0$ by the co-area theorem.

Since p_0 is quasi-homogeneous, I_{x^3} can be also computed by a standard ([1]) change of variables (from $p^{-1}(t)$ to $p^{-1}(1)$, e.g.).

c) For λ arbitrary, we note that (24) and (25) imply $I_{x^3 - \lambda^2 x - 2y^2} = 0$. This also follows from the fact that the form $(x^3 - \lambda^2 x - 2y^2)dx dy/dp$ is exact. Indeed, we have $(x^3 - \lambda^2 x - 2y^2)dx \wedge dy = dp \wedge d\theta$, for $\theta = xy$.

4.2. Let $p = p_u$ be the polynomial defined by

$$p_u(z) = |z^4| - (z^2 + \bar{z}^2) - 2u|z^2|, \quad (26)$$

where $u \in \mathbf{C}$ is a parameter, $z \in \mathbf{C}$ is the variable.

Rewriting $z = x + iy$ gives

$$p_u(x, y) = (x^2 + y^2)^2 - 2(x^2 - y^2) - 2u(x^2 + y^2).$$

Allowing next x, y to be complex, we are in the case $n = 2$.

a) For u real, $u \neq 0$, the domain

$$\Omega_u := \{z \in \mathbf{C} \mid p_u(z) = -(u-1)^2\} \quad (27)$$

is relatively compact in \mathbf{C} and for $u \neq 0$ it is a *quadrature domain* ([2]), i.e.

$$\partial\Omega_u = |P(z)|^2 - \sum_{0 \leq k < d} |Q_k(z, \bar{z})|^2 = 0, \quad (28)$$

where P, Q_k are polynomials, $d = \deg P$ and $\deg Q_k = k$.

Indeed, in the present case, we can write $\partial\Omega_u = \{Q_u(z, \bar{z}) = 0\}$, where

$$Q_u(z, \bar{z}) = |z^2 - 1| - 2u(|z|^2 + 1) + u^2 \quad (29)$$

is of the form (28); $\partial\Omega_u$ is the boundary of a Hele-Shaw flow ([11]) with two sources at ± 1 , which for $0 < u < 1$ is a disjoint union of two disks with centers at ± 1 .

For 1-parameter quadrature domains Ω_u , the moments can be computed from the equation of the boundary $Q_u(z, \bar{z}) = 0$ by ([12]):

$$\frac{-\pi Q'}{Q} = \sum_{k,l=0}^{\infty} \frac{a'_{kl}(u)}{z^{k+1}\bar{z}^{l+1}} \quad (30)$$

where Q' denotes derivative w. r. t. the parameter u . The moments (in the (z, \bar{z}) writing) $a_{kl}(u)$ are defined by

$$a_{kl}(u) := \int \int_{\Omega_u} z^k \bar{z}^l dA,$$

where $dA = dx dy$ is the area element.

Computing the left-hand term of (30) in this case (using $\zeta = 1/z$ and long division), we obtain

$$a'_{00} = 2\pi, a'_{11} = 2\pi(u+1), a'_{20} = a'_{02} = 2\pi$$

(and besides the vanishing of the moments of order one).

In coordinates (x, y) this gives:

$$I_0 = 2\pi, I_{x^2} = \frac{\pi}{2}(2u+3), I_{y^2} = \frac{\pi}{2}(2u+1),$$

the integrals being taken over $\gamma_u(t) := \partial\Omega_u$, where $t = -(u-1)^2$.

b) On the other hand, we will show that the algorithm of section 3 implies

$$-I_{x^2}(t) \cdot (1+u) + I_{y^2}(t) \cdot (1-u) = (t - \frac{1}{2}\partial_t^{-1})I_0 \text{ mod } \mathcal{O}_{u,t} \quad (31)$$

w.r.t. a $\gamma_u(t)$ as in Thm 0.4.1.

Indeed, from $I_p = tI_0$, we have first

$$I_{(x^2+y^2)^2} - 2I_{x^2}(1+u) + 2I_{y^2}(1-u) = tI_0. \quad (32)$$

We have $\partial_x p = 4x(x^2 + y^2 - 1 - u)$, $\partial_y p = 4y(x^2 + y^2 + 1 - u)$. For $u \neq \pm 1$, there are 5 non-degenerate critical points : $0, (\pm\sqrt{1+u}, 0), (0, \pm\sqrt{u-1})$; for $u = \pm 1$, the critical point 0 has Milnor number $\mu = 3$.

By the algorithm of section 3 (parametric version, at a regular value of the parameter) for

$$f = f_u := (x^2 + y^2)^2 - (1+u)x^2 + (1-u)y^2 = (1/4)x\partial_x p + (1/4)y\partial_y p,$$

we have $g^0 = (1/4)(x, y)$, $f_1 = \operatorname{div} g^0 = 1/2$ and hence $I_f = (1/2)\partial_t^{-1}I_0$. Explicitly, this gives

$$I_{(x^2+y^2)^2} - (1+u)I_{x^2} + (1-u)I_{y^2} = \frac{1}{2}\partial_t^{-1}I_0.$$

Subtracting this from (32), we obtain formula (31) of this sub-paragraph.

c) Finally, in order to compare the results at a) and b) above, we take $t = -(u-1)^2$, or equivalently $u = \sqrt{-t} + 1$ in (31) (here $u > 1$ s.t. $u = \text{regular parameter}$; $t < 0$ avoids the critical values of p_u).

We obtain, by b)

$$-I_{x^2}(t)(\sqrt{-t} + 2) + I_{y^2}(t)(-\sqrt{-t}) = (t - \frac{1}{2}\partial_t^{-1})I_0(t) \bmod \mathcal{O}_{\sqrt{-t}}. \quad (33)$$

By a), the right-hand term equals πt ; the left-hand term is in $\mathcal{O}_{\sqrt{-t}}$, since both $I_{x^2}(t)$, $I_{y^2}(t)$ depend linearly in $\sqrt{-t}$. Therefore a) too implies that equality mod $\mathcal{O}_{\sqrt{-t}}$ (and not exact equality) holds in (33).

4.3. a) Let $n \geq 1$ and let $p \in \mathbf{C}[x]$ ($x = (x_1, \dots, x_n)$) be a polynomial satisfying (*1) and such that its critical points $(x_j)_{1 \leq j \leq s}$ are non-degenerate; (note that $s = \mu_T$).

Let f be a polynomial of the form

$$f = x^\beta + h, \quad (34)$$

where $h \in I_{\nabla p}$, $\beta \in I$. For the Fourier integral with phase p and amplitude f defined by (notations differs from 1.3)

$$I_f^\Gamma(\tau) := \int_\Gamma e^{i\tau p(x)} f(x) dx,$$

(where $\Gamma = \Gamma^{(n)} \in H_n^\Phi(\mathbf{C}^n)$, $\Phi = \Phi(\theta)$, θ a generic direction in \mathbf{C}) we have, by the first step of the algorithm of Section 3,

$$I_f^\gamma(\tau) = I_{x^\beta}^\Gamma(\tau) + \frac{1}{i\tau} I_{f_1}^\Gamma(\tau),$$

for some $f_1 \in \mathbf{C}[x]$.

b) We shall use the method of the stationary phase to show that

$$I_f^\Gamma(\tau) \sim I_{x^\beta}^\Gamma(\tau) \cdot (1 + O(\frac{1}{\tau})), \quad (35)$$

asymptotically as $\tau \rightarrow \infty$, τ in a sector in \mathbf{C} . (This checks asymptotically the first step of the algorithm of Thm. 0.4.1).

Since (1.2) $\Gamma = \sum_{1 \leq j \leq \mu} \Gamma_j$, it is sufficient to prove the above formula with Γ replaced by Γ_j , in local coordinates x' at x_j . Let $\Gamma_j = m_{\Gamma_j} \cdot \Delta$, where (notations of Section 1) $\Delta \in H_n(S_j, S_j^+)$ is the standard (Lefschetz) generator (x_j is non-degenerate).

By stationary phase (since $e^{i\tau p(x_j)} = (1 + O(\frac{1}{\tau}))$), we have (asymptotically as $\tau \rightarrow \infty$, τ in a sector in \mathbf{C})

$$I_f^{\Gamma_j}(\tau) \sim \frac{1}{\tau^{n/2}} (1 + O(\frac{1}{\tau})) \cdot C_j \cdot K_n,$$

with constants C_j, K_n defined by

$$C_j = C_j(f, \Gamma_j, p) = f(x_j) \cdot (\text{Hess } p(x_j))^{-1/2} \cdot m_{\Gamma_j},$$

$$K_n = (2\pi)^{n/2} \cdot e^{i\pi/4}.$$

Since f was assumed of the form (34), writing $f(x) = x^\beta + \sum_{1 \leq k \leq n} g_k(x) \partial_k p(x)$, we note that $f(x_j) = x_j^\beta$. Therefore

$$C_j(f, \Gamma_j, p) = C_j(x^\beta, \Gamma_j, p),$$

and from this (by a similar computation for the right hand term of (35)) we obtain the asymptotic equality of (35).

c) In particular, for $n = 2$, let

$$p(x, y) = \frac{x^{\mu+1}}{\mu+1} - x + y^2.$$

Since $\nabla p = (x^\mu - 1, 2y)$, the critical points of p are $(\epsilon^j)_{0 \leq j \leq \mu-1}$, where ϵ is a primitive root of order μ of 1.

Letting $f := x^{\mu+1} = x + x(x^\mu - 1)$, $\mu \geq 2$, we are in the situation above. By the algorithm of Section 3, if $[\gamma(t)]$ is parallel transported in $\mathbb{H}_{n-1}(p^{-1}(t))$ (t large in a sector of axis θ), we have

$$I_{x^{\mu+1}}^\gamma = I_x^\gamma + \partial_t^{-1} I_0^\gamma \text{ mod } \mathcal{O}_t.$$

(For $\mu = 2$, we have used this in example 4.1).

By the above computation using the stationary phase formula (for Γ s.t. $\partial\Gamma = \gamma(t_0)$, some t_0), the leading terms as $\tau \rightarrow \infty$ of $I_{x^{\mu+1}}^\Gamma$ and I_x^Γ are both equal with

$$\frac{1}{\tau} \pi \cdot e^{i\pi/4} \sqrt{\frac{2}{\mu}} \sum_{0 \leq j \leq \mu-1} \epsilon^{j/2} \cdot m_{\Gamma_j}.$$

d) For $\mu = 2$, $n = 1$, we also have (similarly) the estimate (for a Laplace transform with phase)

$$\int_{\Gamma} e^{-\tau(\frac{x^3}{3}-x)} dx \sim i \sqrt{\frac{\pi}{\tau}} \cdot (1 + O(\frac{1}{\tau})),$$

which is used in computations for asymptotics ([4],[18]) of the Airy function (where directionally only one of the critical points x_j contributes and $\Gamma = \Gamma_j$ is s.t. $m_{\Gamma_j} = 1$).

REFERENCES

- [1] Arnold, V.I.; Gusein-Zade, S.M.; Varchenko, A.N.: *Singularities of differentiable maps II*. Birkhauser 1988.
- [2] Aharonov, D; Shapiro, H.S.: Domains on which analytic functions satisfy quadrature identities. *J. d'analyse mathematique* 30,p.39-73,1976.
- [3] Banica, C.; Stanasila, O.: *Methodes algebriques dans la theorie globale des espaces complexes I*. Varia math, Gauthier-Villars 1977.
- [4] Bleistein, N.; Handelsman, R.A.: *Asymptotic expansions of integrals*. Dover 1986.
- [5] Cassier, G.: Probleme des moments sur un compact de \mathbf{R}^n . *J. functional analysis* 58,p.254-266,1984.
- [6] Cox, D.; Little, J.; O'Shea, D.: *Ideals, varieties and algorithms*. 2nd ed., Springer 1997.
- [7] Fuglede, B.: The multidimensional moment problem. *Expo. math.* 1,p.47-65,1983.
- [8] Goursat, E.: *A course in mathematical analysis I*. Dover 1959.
- [9] Grothendieck, A.: *Seminaire de Geometrie Algebrique II*. IHES.
- [10] Greuel, G., Pfister, G.: *A singular introduction to commutative algebra*. Springer 2002.
- [11] Gustaffsson, B.: Singularities and special points on a quadrature domain from an algebraic point of view. *J. d'analyse math.* 51,p.91-117,1988.
- [12] Gustaffsson, B.; Putinar, M.: The exponential transform: a renormalized Riesz potential at a critical point. *Ind. Univ. Math. J.* 52(3),p.527-568,2003.

- [13] Landau, H., ed.: *Moments in mathematics*. Proc. symp. appl. math. 37, AMS, 1987.
- [14] Maisonobe, Ph.; Rombaldi, J.E.: Solutions du systeme de Gauss-Manin d'un germe de fonction a point critique isole. *Singularites des systemes differentiels de Gauss-Manin*, Progr. Math. 2,p.307-339,Birkhauser 1979.
- [15] Malgrange, B.: Integrales asymptotiques et monodromie. *Ann. scient. Ec. Norm. Sup.* 4^e serie,t.7,p.405-430,1974.
- [16] Malgrange, B.: Sur les points singuliers des equations differentielles. *L'enseignement math.*,XX(3-4),1974.
- [17] Malgrange, B.: Equations differentielles a coefficients polynomiaux. *Progr. math.* 96, Birkhauser 1991.
- [18] Murray, J.D.: *Asymptotic analysis*. Clarendon Press, Oxford 1974.
- [19] Pham, F.: *Singularites des systemes differentiels de Gauss-Manin*, Progr. Math. 2, p.1-24, Birkhauser 1979.
- [20] Pham, F.: La descente des cols par les onglets de Lefschetz avec vue sur Gauss-Manin. *Asterisque* 130,p.11-47,1985.
- [21] Pham, F.: Transformees de Laplace des microsolutions de systemes holonomes. *L'enseignement math.* 30,p.57-84,1984.
- [22] Putinar, G.: Asymptotics for extremal moments and monodromy of complex singularities. *submitted*.
- [23] Putinar, M.: Extremal solutions of the two-dimensional L-problem of moments I, *J.F.A.*, 136, p. 331-364, 1996; II, *J. approx. theory*, 92, no.1,p.38-58, 1998.
- [24] Putinar, G.; Putinar, M.: Reconstruction of algebraic sets from dynamic moments. *submitted*.
