

Final Exam
Math 240 B
Winter 2011
Prof. Ye

Your name
Your perm. number

Scores

- 1.
- 2.
- 3.
- 4.
- 5.
- 6.
- 7.

Total:

Each problem is worth 16 points. In particular, 2 extra credit points are included.

**Please email your final and homework to Prof. Ye:
yer@math.ucsb.edu**

**You can type up or scan your documents. The final is due on Saturday,
March 19, 2010. Please keep your originals.**

1. Let M be a smooth manifold and $\alpha \in \Omega^m(M)$, i. e. α is a smooth form of degree m on M .

1) Assume $m = 1$. Prove the following formula for the exterior derivative

$$d\alpha(X, Y) = X\alpha(Y) - Y\alpha(X) - \alpha([X, Y]) \quad (0.1)$$

for smooth vector fields X and Y .

2) Prove the following general formula for the exterior derivative

$$\begin{aligned} d\alpha(X_1, \dots, X_{m+1}) &= \sum_{i=1}^{m+1} (-1)^{i-1} X_i \alpha(X_1, \dots, \hat{X}_i, \dots, X_{m+1}) \\ &\quad + \sum_{i < j} (-1)^{i+j} \alpha([X_i, X_j], X_1, \dots, \hat{X}_i, \dots, \hat{X}_j, \dots, X_{m+1}) \end{aligned}$$

for arbitrary smooth vector fields X_i , where the hat means that the term is omitted.

Note: In John Lee's book, a proof of this formula is given. You are required to find a proof without reading Lee's proof, until you have become completely frustrated (if that happens at all).

2. Consider a smooth manifold M of dimension n . Let $\Omega^*(M) = \bigoplus_{0 \leq k \leq n} \Omega^k(M)$ denote the algebra of smooth differential forms on M . Let $\Omega_c^*(M) = \bigoplus_{0 \leq k \leq n} \Omega_c^k(M)$ denote the subalgebra of $\Omega(M)$ consisting of smooth differential forms on M with compact support.

1) Show that the De Rham cohomology $H_{dR,c}^*(M)$ with compact support can be defined analogously to $H_{dR}^*(M)$.

2) Show that the inclusion map of $\Omega_c^*(M)$ into $\Omega^*(M)$ induces a natural homomorphism $F : H_{dR,c}^*(M) \rightarrow H_{dR}^*(M)$.

3) Show that F is not injective for $M = \mathbf{R}$ (the real line).

4) Assume that M is noncompact. Show that the restriction of F to $H_{dR,c}^k(M)$ is not injective for each $0 < k \leq n$. Hint: Use Stokes Theorem.

5) Assume that M is connected and noncompact. Show that $H_{dR,c}^0(M) = \{0\}$.

3. 1) Show that the 1-dimensional De Rham cohomology group $H_{dR}^1(S^1 \times S^1)$ is isomorphic to \mathbf{R}^2 .

2) Let $n \geq 2$ be general. Show that the 1-dimensional De Rham cohomology group $H_{dR}^1(T^n)$ of the n -dimensional torus $T^n = S^1 \times \dots \times S^1$ (n factors) is isomorphic to \mathbf{R}^n

Hint: Extend the proof for the case $n = 1$ presented in the lectures.

4. Let ∇ be a connection on the tangent bundle of a smooth manifold M . Recall that it extends to all tensor fields, i. e. we can take covariant derivatives $\nabla_v \sigma$ for smooth tensor fields σ . Let $\gamma : [a, b] \rightarrow M$ be a smooth curve, and $\sigma = \sigma(t)$ a smooth tensor

field of some type (r, s) along γ , i. e. $\sigma(t) \in T_s^r(T_{\gamma(t)}M)$ for each t .

1) We define the covariant derivative $\nabla_{\frac{d}{dt}}\sigma$ as follows. Let $[a', b']$ be a subinterval of $[a, b]$ such that $\gamma([a', b'])$ is contained in a coordinate chart (U, ϕ) . Let $\sigma_{j_1 \dots j_s}^{i_1 \dots i_r}$ be basis tensor fields in the chart, i. e.

$$\sigma_{j_1 \dots j_s}^{i_1 \dots i_r} = dx^{i_1} \otimes \dots \otimes dx^{i_r} \otimes \frac{\partial}{\partial x^{j_1}} \otimes \dots \otimes \frac{\partial}{\partial x^{j_s}}. \quad (0.2)$$

We have

$$\sigma(t) = \sum a_{i_1 \dots i_r}^{j_1 \dots j_s}(t) \sigma_{j_1 \dots j_s}^{i_1 \dots i_r}|_{\gamma(t)} \quad (0.3)$$

for some smooth functions $a_{i_1 \dots i_r}^{j_1 \dots j_s}(t)$. Then we set for $a' \leq t \leq b'$

$$\nabla_{\frac{d}{dt}}\sigma = \sum \left(\frac{d}{dt} a_{i_1 \dots i_r}^{j_1 \dots j_s}(t) \right) \sigma_{j_1 \dots j_s}^{i_1 \dots i_r}|_{\gamma(t)} + \sum a_{i_1 \dots i_r}^{j_1 \dots j_s}(t) \nabla_{\gamma'(t)} \sigma_{j_1 \dots j_s}^{i_1 \dots i_r}. \quad (0.4)$$

Show that this definition is independent of the choice of the chart.

2) Let $\sigma_0 \in T_s^r(T_{\gamma(a)}M)$ for given r and s . Show that there is a *unique* parallel smooth tensor field σ of type (r, s) along γ , such that $\sigma(0) = \sigma_0$. Here “parallel” means $\nabla_{\frac{d}{dt}}\sigma \equiv 0$. ($\sigma(t)$ is called the parallel transport of σ_0 along the curve γ .)

Hint: First handle the case of type $(0, 1)$, i. e. the case of vector fields. This special case is worth more than half of the credit.

5. Let (M, g, ∇) be a Riemannian manifold together with the Levi-Civita connection of g .

1) Let $\gamma : [a, b] \rightarrow M$ be a smooth curve and $\{v_1, \dots, v_n\}$ an orthonormal basis of the tangent space at $\gamma(a)$. Let $e_i(t)$ be the parallel transport of v_i along γ . Show that $\{e_i(t)\}$ is orthonormal for each t .

2) Assume in addition that M is oriented and (v_1, \dots, v_n) is a positive basis. Show that $(e_1(t), \dots, e_n(t))$ is a positive basis for each t .

6. Let (M, g, ∇) be an oriented Riemannian manifold together with the Levi-Civita connection of g .

1) Show that g is parallel, i. e. $\nabla_v g = 0$ at each point $p \in M$ and for each tangent vector $v \in T_p M$.

2) Show that the volume form $dvol$ is parallel, i. e. $\nabla_v dvol = 0$ at each point $p \in M$ and for each tangent vector $v \in T_p M$.

7. Assume the following result: $H_{dR}^n(M)$ is isomorphic to \mathbf{R} for a connect, closed and orientable smooth manifold M of dimension n . Prove that $H_{dR}^n(M) = \{0\}$ for a connected, closed and non-orientable manifold M of dimension n . Hint: utilize the oriented double cover of M .